

Composition of Axial Functions of Product of Finite Sets

Krzysztof Płotka
Department of Mathematics
University of Scranton
Scranton, PA 18510, USA
plotkak2@scranton.edu

September 14, 2004

To the memory of my Father

Abstract

We show that every function $f: A \times B \rightarrow A \times B$, where $|A| \leq 3$ and $|B| < \omega$, can be represented as a composition $f_1 \circ f_2 \circ f_3 \circ f_4$ of four axial functions, where f_1 is a vertical function. We also prove that for every finite set A of cardinality at least 3, there exist a finite set B and a function $f: A \times B \rightarrow A \times B$ such that $f \neq f_1 \circ f_2 \circ f_3 \circ f_4$, for any axial functions f_1, f_2, f_3, f_4 , whenever f_1 is a horizontal function.

A function $f: A \times B \rightarrow A \times B$ is called *vertical* ($f \in V$) if there exist a function $f_1: A \times B \rightarrow A$ such that $f(a, b) = (f_1(a, b), b)$. It is called *horizontal* ($f \in H$) if $f(a, b) = (a, f_2(a, b))$ for some function $f_2: A \times B \rightarrow B$. If f is horizontal or vertical then we call it *axial*. A one-to-one function from $A \times B$ onto $A \times B$ is called a *permutation* of $A \times B$. The family of all functions from $A \times B$ into $A \times B$ is denoted by $(A \times B)^{A \times B}$. If $F_1, F_2, \dots, F_n \subseteq (A \times B)^{A \times B}$, then we write $F_1 F_2 \dots F_n$ to denote $\{f_1 \circ f_2 \circ \dots \circ f_n : f_i \in F_i, i = 1, 2, \dots, n\}$.

AMS Subject Classification (2000): Primary **03E20**; Secondary: 08A02.

Key words: axial functions.

It is convenient, especially when the set $A \times B$ is finite, to use matrices to represent functions from $(A \times B)^{A \times B}$. Given a function $f \in (A \times B)^{A \times B}$ and a matrix $M = [m_{(a,b)}]$ of size $|A| \times |B|$, we define $f[M] = [m_{f(a,b)}]$. If the elements of the matrix $M = [m_{(a,b)}]$ are distinct, then the matrix $f[M]$ uniquely determines the function f . We will often identify the elements (a, b) of $A \times B$ and the corresponding entries $m_{(a,b)}$ of the matrix M . Observe also that if $f, g \in (A \times B)^{A \times B}$, then $f[g[M]] = (g \circ f)[M]$. To see this let $M' = g[M]$ and $M'' = f[M'] = f[g[M]]$. For $(a, b) \in A \times B$ we have that $m''_{(a,b)} = m'_{(c,d)}$ for some $(c, d) \in A \times B$ such that $(c, d) = f(a, b)$. Now, note that $m'_{(c,d)} = m_{g(c,d)}$. Hence $m''_{(a,b)} = m_{g(c,d)} = m_{g(f(a,b))} = m_{(g \circ f)(a,b)}$.

Banach ([M], Problem 47) asked whether every permutation function of a cartesian product of two infinite countable sets can be represented as a composition of finitely many axial functions. The question was answered affirmatively by Nosarzewska [N]. Further research on this subject was done by Ehrenfeucht and Grzegorek [EG, G]. They discussed the smallest possible number of axial functions needed. Also, they considered the case when both sets are finite. In particular, they proved the following.

Theorem 1. *Let $f, p \in (A \times B)^{A \times B}$ and p be a permutation.*

- (i) *Then $p = p_1 \circ p_2 \circ p_3 \circ p_4$, where all p_i are axial permutations of $A \times B$.*
- (ii) *If A is finite, then $p = p_1 \circ p_2 \circ p_3$, where all p_i are axial permutations of $A \times B$ and $p_1 \in V$.*
- (iii) *If $A \times B$ is infinite, then $f = f_1 \circ f_2 \circ f_3$, where all f_i are axial functions.*
- (iv) *If $A \times B$ is finite, then f can be represented as $f = f_1 \circ \dots \circ f_6$, where all f_i are axial functions and $f_1 \in V$.*

Let us mention here, that Ehrenfeucht and Grzegorek showed also that it is not possible to decrease the numbers 4 in part (i) and 3 in parts (ii) and (iii) of the above theorem. In addition, in part (i) it cannot be specified that $p_1 \in V$. However, it was not proved that 6 in part (iii) is the smallest possible. Later, Szyszkowski [S] proved that 6 can be decreased to 5. He also gave an example which showed that 6 cannot be decreased to 3. In his

example one of the sets has at least 4 elements and the other one at least 5. Below, we present an example in which both sets have exactly three elements each. It is worthy to mention here that, if one of the sets has at most two elements, then 3 axial functions are enough (see Remark 5).

Example 2 *The number 6 in Theorem 1 (iv) cannot be reduced to 3.*

$$\text{Let } M = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}. \text{ Define } f \in (A \times B)^{A \times B} \text{ by } f[M] = \begin{bmatrix} i & g & e \\ d & h & b \\ h & a & c \end{bmatrix}.$$

We will justify why $f \neq f_1 \circ f_2 \circ f_3$ for any axial functions f_1, f_2, f_3 such that $f_1 \in V$ (the case when $f_1 \in H$ is analogous). Notice that for the equality $f[M] = (f_1 \circ f_2 \circ f_3)[M] = f_3[f_2[f_1[M]]]$ to hold, every column of $f[M]$ would have to be contained in a selector from the rows of $f_1[M]$. In particular, since the sets $\{i, d, h\}$ and $\{g, h, a\}$ form two columns of $f[M]$, the elements a, d, g would have to appear in different rows of $f_1[M]$ than the element h . But this is impossible. ■

The question, which remains open, is whether every function can be obtained as a composition of four axial functions. We give partial answers to this question. The following holds.

Theorem 3.

- (i) *If $|A| = 3$, then $(A \times B)^{A \times B} = \text{VHVH}$.*
- (ii) *If $|A| \geq 3$, then there exists an integer m_0 such that $(A \times B)^{A \times B} \neq \text{HVHV}$, whenever $|B| \geq m_0$.*

Theorem 3 implies the following.

Corollary 4 *There exist finite sets A, B and a function $f: A \times B \rightarrow A \times B$ such that $f \in \text{VHVH}$ and $f \notin \text{HVHV}$.*

PROOF OF THEOREM 3. (i) First observe that it suffices to prove the result for functions $f: A \times B \rightarrow A \times B$ such that the entries in each row of the matrix $f[M]$ are all distinct, that is, $|f(\{a\} \times B)| = |B|$ for every $a \in A$. This

is so, because for an arbitrary function $f': A \times B \rightarrow A \times B$, there exists a function f of the above type and a horizontal function $h: A \times B \rightarrow A \times B$ such that $f'[M] = h[f[M]] = (f \circ h)[M]$. Since the composition of two horizontal functions is a horizontal function, if $f \in \text{VHVH}$, then also $f' \in \text{VHVH}$. Hence, let f be a function such that the entries in each row of the matrix $f[M]$ are all distinct. It can be easily proved by induction on $|B|$, that there exists a horizontal permutation $h: A \times B \rightarrow A \times B$ such that there exists a partition of B into three sets B_1, B_2, B_3 (some of the sets may be empty) with the following properties:

1. $|(f \circ h)(A \times \{b_1\})| = 1$ for every $b_1 \in B_1$,
2. $|(f \circ h)(A \times \{b_2\})| = 2$ for every $b_2 \in B_2$,
3. $|(f \circ h)^{-1}(\{m\})| \leq 2$ for every $m \in (f \circ h)(A \times B_3)$ and $|\{m \in (f \circ h)(A \times B_3) : |(f \circ h)^{-1}(\{m\})| = 2\}| \equiv 0 \pmod{3}$,
4. if $(f \circ h)(A \times \{b\}) \cap (f \circ h)(A \times \{b'\}) \neq \emptyset$, then $b, b' \in B_3$.

For the matrix $h[f[M]]$ this means that each column with index in B_1 has all the entries equal, each column with index in B_2 has only two different entries, and the number of entries appearing twice in the part of $h[f[M]]$ corresponding to B_3 is divisible by 3. In addition, columns with indices in B_3 are the only columns which can share entries with other columns.

$$h[f[M]] = \underbrace{\begin{bmatrix} a & b & \dots & p & q & \dots & v & x & y & \dots \\ a & b & \dots & p & s & \dots & v & w & z & \dots \\ a & b & \dots & r & s & \dots & w & x & z & \dots \end{bmatrix}}_{\substack{B_1 \quad B_2 \quad B_3}}$$

So, if we can prove that $f \circ h \in \text{VHVH}$, then, since h is a horizontal permutation, we will also prove that $f \in \text{VHVH}$. Hence, without loss of generality, we can assume that f itself satisfies the four above conditions. Now, let us partition the set $(A \times B) \setminus f(A \times B)$ into sets E_1, E_2, E_3 such that $|E_1| = 2|B_1|$ and $|E_2| = |B_2|$. Next define the partition $\mathcal{P} = \mathcal{P}_1 \cup \mathcal{P}_2 \cup \mathcal{P}_3 \cup \mathcal{P}_4 \cup \mathcal{P}_5$ of $A \times B$ into sets of size 3 as follows.

- $\{m_1, m_2, m_3\} \in \mathcal{P}_1$ if $m_1 \in f(A \times B_1)$, $m_2, m_3 \in E_1$

- $\{m_1, m_2, m_3\} \in \mathcal{P}_2$ if $m_1, m_2 \in f(A \times \{b_2\})$ for some $b_2 \in B_2, m_3 \in E_2$
- $\{m_1, m_2, m_3\} \in \mathcal{P}_3$ if $m_1, m_2, m_3 \in E_3$
- $\{m_1, m_2, m_3\} \in \mathcal{P}_4$ if $m_i \in f(A \times B_3), |f^{-1}(\{m_i\})| = 2$ for $i = 1, 2, 3$
- $\{m_1, m_2, m_3\} \in \mathcal{P}_5$ if $m_i \in f(A \times B_3), |f^{-1}(\{m_i\})| = 1$ for $i = 1, 2, 3$

The existence of this partition follows from the conditions 1-4. Note that these conditions also imply that $|\mathcal{P}_3| = |\mathcal{P}_4|$.

Based on Theorem 1 (ii), there exists a vertical permutation $f_1 \in V$ such that for each set $P \in \mathcal{P}$, the elements of P appear in different rows of $f_1[M]$ (see the argument in Example 2). Now observe that there exists a horizontal permutation $f_2 \in H$ such that the columns of $f_2[f_1[M]]$ are the sets of the partition \mathcal{P} and the sets from $\mathcal{P}_1 \cup \mathcal{P}_2$ correspond to columns with indices in $B_1 \cup B_2$. The function f_2 can be modified, so the elements from the sets in \mathcal{P}_3 are replaced by the elements from the sets in \mathcal{P}_4 and the latter appear twice in the matrix $f_2[f_1[M]]$. Notice that the parts of matrices $f[M]$ and $f_2[f_1[M]]$ corresponding to $A \times B_3$ are permutations of each other, so by Theorem 1 (ii) one can be obtained from the other by performing three axial permutations (note that if both sets A and B are finite, then Theorem 1 (ii) with $p_1 \in V$ replaced by $p_1 \in H$ also holds). Additionally, the columns in $f_2[f_1[M]]$ with indices in $B_1 \cup B_2$ can be made identical to the columns of $f[M]$ by performing one vertical operation. Consequently, we can claim that there exist three axial functions $f_3 \in H, f_4 \in V, f_5 \in H$ such that $f[M] = f_5[f_4[f_3[f_2[f_1[M]]]]$. Hence $f = f_1 \circ \dots \circ f_5$. Since $f_2 \in H$ and $f_3 \in H$, we obtain that $f_2 \circ f_3 \in H$ and finally $f \in VHVH$.

(ii) Denote $|A|$ by k . Define $n = k(k-1) + 1$ and $m_0 = \left\lceil \frac{k\binom{n}{k} - n}{k-2} \right\rceil$. Let $m \geq m_0$ be an integer and M be a $k \times m$ matrix with all entries distinct and such that the first n entries in the first row are a_1, a_2, \dots, a_n .

We define a function $f: A \times B \rightarrow A \times B$ ($|B| = m$) by defining $f[M]$. The entries from the “bottom” $(k-2)$ rows of M do not appear in $f[M]$. The first $\binom{n}{|A|} = \binom{n}{k}$ columns of $f[M]$ form all k -subsets of $\{a_1, a_2, \dots, a_n\}$. The rest $(m - \binom{n}{k})$ of the columns of $f[M]$ are formed using all the entries from the first two rows of M except a_1, a_2, \dots, a_n (some of them may need to appear

more than once). Note that this is possible because the number $(2m - n)$ of entries in the first two rows of M except a_1, a_2, \dots, a_n is not greater than the number $(k(m - \binom{n}{k}))$ of positions in $m - \binom{n}{k}$ columns of $f[M]$. Indeed, $m \geq m_0 = \left\lceil \frac{k\binom{n}{k} - n}{k-2} \right\rceil \geq \frac{k\binom{n}{k} - n}{k-2}$. Consequently, $m(k-2) \geq k\binom{n}{k} - n$ and $k(m - \binom{n}{k}) \geq 2m - n$.

$$M = \underbrace{\begin{bmatrix} a_1 & a_2 & a_3 & \dots & & a_n & \dots \\ \cdot & \cdot & \cdot & & & & \\ \vdots & \vdots & \vdots & & & & \vdots \\ \cdot & \cdot & \cdot & & & & \end{bmatrix}}_{m \text{ columns}}$$

$$f[M] = \underbrace{\begin{bmatrix} a_{i_1} & a_{j_1} & \dots & & \dots \\ a_{i_2} & a_{j_2} & \dots & & \dots \\ \vdots & \vdots & \vdots & & \vdots \\ a_{i_k} & a_{j_k} & \dots & & \dots \end{bmatrix}}_{\substack{\binom{n}{k} \text{ columns} \\ \text{all } k\text{-subsets of} \\ \{a_1, a_2, \dots, a_n\}}} \underbrace{\hspace{10em}}_{\substack{m - \binom{n}{k} \text{ columns} \\ \text{all entries from the first two} \\ \text{rows of } M \text{ except } a_1, a_2, \dots, a_n}}$$

We will show that $f \notin \text{HVHV}$. Assume, by the way of contradiction, that there exist axial functions f_1, \dots, f_4 such that $f_1 \in \text{H}$ and $f = f_1 \circ f_2 \circ f_3 \circ f_4$, i.e. $f[M] = f_4[f_3[f_2[f_1[M]]]]$.

Claim 1. There are k elements out of $\{a_1, a_2, \dots, a_n\}$, say a_1, a_2, \dots, a_k , such that there is a row of the matrix $f_2[f_1[M]]$ which does not contain any of these elements. To see this, first observe that f_1 restricted to the first two rows of M is a permutation. This is so, because all the elements of M from the first two rows appear in the final matrix $f[M]$. Let us call the elements from the second row of $f_1[M]$ that appear in the same columns as a_1, a_2, \dots, a_n by b_1, b_2, \dots, b_n , respectively. Since $n = k(k-1)$ and $|A| = k$, by the Pigeonhole Principle, there exists a row in $f_2[f_1[M]]$ which contains at least k elements out of b_1, b_2, \dots, b_n . This row does not contain at least k elements out of a_1, a_2, \dots, a_n .

Claim 2. Applying the functions $f_3 \in \text{H}$ and $f_4 \in \text{V}$ to the matrix $f_2[f_1[M]]$ will not result in a matrix containing a column whose entries are

a_1, a_2, \dots, a_k . Hence $f[M] \neq f_4[f_3[f_2[f_1[M]]]]$. A contradiction. ■

Let us mention here that, using similar technique to the one used in the proof of Theorem 3 (i), we can show the following.

Remark 5 Let $|A| = 2$. Then $(A \times B)^{A \times B} = \text{HVH}$. In addition, if $|B| \geq 3$, then $(A \times B)^{A \times B} \neq \text{VHV}$.

The counterexample justifying the second part of the remark can be found in [S] (page 36).

References

- [EG] A. Ehrenfeucht, E. Grzegorek, *On axial maps of direct products I*, Colloq. Math. **32** (1974), 1–11.
- [G] E. Grzegorek, *On axial maps of direct products II*, Colloq. Math. **34** (1976), 145–164.
- [M] R. D. Mauldin (Ed.), *The Scottish Book; Math at the Scottish Cafe*, Birkhäuser, Boston, MA, 1981.
- [N] M. Nosarzewska, *On a Banach's problem of infinite matrices* Colloq. Math. **2** (1951), 194–197.
- [S] M. Szyszkowski, *On axial maps of the direct product of finite sets*, Colloq. Math. **75** (1998), 33–37.